

SEAL

DO NOT OPEN THE SEAL UNTIL INSTRUCTED TO DO SO

Series :

A

Question Booklet No.

102578

GM/A/07/2024 EXAM.

2024

PAPER—II

Invigilator's Signature

Candidate's Signature

Time : 2 Hours

Maximum Marks : 200

ROLL NO.

--	--	--	--	--	--

INSTRUCTIONS FOR CANDIDATES

1. Immediately after the commencement of the Examination, candidate should check that this Booklet does **NOT** have any unprinted, torn or missing pages/Sl. No. etc. If any defect is found, candidates should not write or mark anything on the OMR RESPONSE SHEET and immediately report it to the room Invigilator for replacement by a Complete Question Booklet.
2. Candidate should carefully read the instructions on the back of the OMR RESPONSE SHEET. They should **NOT** write Name, mark, make any stray marking or write anything irrelevant on either side of the OMR RESPONSE SHEET. Remarks other than the Answers and requisite details will be treated as revealing your identity and upon physical verification if such remarks are found, the OMR RESPONSE SHEET will be invalidated and the candidature cancelled. No rough work should be done on the OMR RESPONSE SHEET. Rough work space provided in the Question Booklet can be used for the same.
3. Candidate should enter correct and complete digits of his/her Roll Number, Booklet Number and other details in the appropriate boxes and darken the corresponding bubbles in the OMR RESPONSE SHEET.
4. Candidate should **NOT** handle his/her OMR RESPONSE SHEET in such a manner as to mutilate, fold etc.
5. This Question Booklet contains **100** questions carrying 2 (two) marks each. Each question contains four responses. **Only one response/answer** for each question should be marked appropriately in the bubble on the OMR RESPONSE SHEET. If more than one response is marked, the answer will be considered wrong.
6. **Candidates are strictly prohibited to possess any book, notebook or loose paper, calculator, mobile phone, any electronic gadget, digital wristwatch etc., inside the Examination Hall, except his/her Unique ID, Admit Card and writing materials only.**
7. Immediately after the final bell indicating the closure of the Examination, candidates should stop marking answers. Candidates should remain seated till the collection of OMR RESPONSE SHEET by the Invigilator. They will leave the Examination Hall after submission of OMR only after they are permitted by the Invigilator.
8. Violation of any of the above Rules will render the candidate liable to be expelled and disqualified from the Examination and according to the nature and gravity of his/her offence, he/she may be debarred from future Examinations and interviews to be conducted by the Commission and other such organizations.

NB : CANDIDATES ARE ALLOWED TO TAKE THIS QUESTION BOOKLET WITH THEM ONLY AFTER COMPLETION OF 2 (TWO) HOURS OF EXAMINATION TIME.

DO NOT OPEN THE SEAL UNTIL INSTRUCTED TO DO SO

GM/A/07/2024 EXAM. 2024/2-A

SEAL

SEAL

1. Which of the following is the most prevalent species of *Plasmodium* causing malaria in the North-East region of India?

- [A] *Plasmodium vivax*
- [B] *Plasmodium malariae*
- [C] *Plasmodium ovale*
- [D] *Plasmodium falciparum*

2. The North-East region of India has a higher prevalence of HIV due to which of the following factors?

- [A] High rates of intravenous drug use
- [B] Unprotected heterosexual transmission
- [C] High number of blood transfusions
- [D] High levels of mother-to-child transmission

3. Which State in the North-East region of India reports the highest incidence of Japanese Encephalitis?

- [A] Assam
- [B] Nagaland
- [C] Manipur
- [D] Tripura



4. Which population group in the North-East region of India is most vulnerable to tuberculosis?

- [A] Tribal populations
- [B] Urban dwellers
- [C] Children under 5 years
- [D] Pregnant women

5. Which substance is most commonly abused in the North-East region of India, contributing to public health challenges?

- [A] Heroin
- [B] Alcohol
- [C] Cannabis
- [D] Opium

6. In the North-East region of India, harm reduction strategies for HIV include all of the following, **except**

- [A] needle exchange programs
- [B] methadone maintenance therapy
- [C] mass screening campaigns
- [D] safe injection sites

7. Which strategy is vital for controlling Japanese Encephalitis in North-East India?

- [A] Annual vaccination campaigns
- [B] Surveillance of migratory bird populations
- [C] Use of insect repellents
- [D] Implementation of water sanitation programs

8. The incubation period for *Plasmodium falciparum* malaria is generally

- [A] 7–10 days
- [B] 14–21 days
- [C] 21–30 days
- [D] 30–40 days

9. The window period for HIV infection refers to the time between
- [A] exposure to HIV and the appearance of symptoms
 - [B] exposure to HIV and the production of detectable antibodies
 - [C] onset of symptoms and death
 - [D] diagnosis and treatment
10. Employees' State Insurance (ESI) scheme is an outcome of
- [A] Bohr Committee
 - [B] Adarkar Committee
 - [C] Srivastava Committee
 - [D] None of the above
11. What is serial interval?
- [A] Time interval between invasion of infection and appearance of first sign and symptom
 - [B] Time between onset of primary case and secondary case
 - [C] Interval of time between receipt of infection by a host and maximum infectivity
 - [D] Time in which infectious agent is transferred from one host to another
12. If a researcher wants to check for incidence of deafness caused by noise effects in traffic police workers after one year, the study conducted will be
- [A] case-control study
 - [B] cross-sectional study
 - [C] ecological study
 - [D] cohort study

13. Which is the best method to reduce the selection error for a clinical trial conducted among two independent groups?
- [A] Matching
 - [B] Blinding
 - [C] Stratification
 - [D] Randomization
14. As the risk for lung cancer is increasing in the North-East region of India, the Government has constituted a board to formulate the Cigarettes and Other Tobacco Products Act (COTPA). This comes under which level of prevention?
- [A] Primordial
 - [B] Primary
 - [C] Secondary
 - [D] Tertiary
15. A researcher plans to conduct a nutrition survey in North-East India. Which of the following is the most suited to find the variations in dietary plan based on religion?
- [A] Simple sampling
 - [B] Snowball sampling
 - [C] Cluster sampling
 - [D] Stratified sampling
16. A woman is diagnosed to be positive for HIV in her first trimester. What is the appropriate management?
- [A] Start ART immediately and continue till six weeks after delivery
 - [B] Start ART immediately and continue life-long
 - [C] Start ART after delivery and continue life-long
 - [D] Start ART after delivery and continue till six weeks after delivery



17. The primary treatment for Japanese Encephalitis is
- [A] antiviral drugs
 - [B] supportive care
 - [C] antibiotics
 - [D] immunoglobulin therapy
18. BCG vaccination provides protection against
- [A] all forms of tuberculosis
 - [B] pulmonary tuberculosis only
 - [C] severe forms of tuberculosis in children
 - [D] drug-resistant tuberculosis
19. The incidence of a disease is defined as
- [A] the number of new cases occurring in a defined population during a specified time period
 - [B] the total number of cases present in a population at a specific time
 - [C] the number of deaths due to the disease
 - [D] the proportion of the population affected
20. Which of the following is a measure of central tendency in a dataset?
- [A] Incidence
 - [B] Mean
 - [C] Prevalence
 - [D] Relative risk

21. The basic reproductive number (R_0) in epidemiology refers to
- [A] the number of secondary cases generated from one primary case in a fully susceptible population
 - [B] the average number of cases in an epidemic
 - [C] the number of cases needed to establish an outbreak
 - [D] the average number of deaths in a pandemic
22. Which of the following is **not** an impact of deforestation on public health?
- [A] Increased incidence of vector-borne diseases
 - [B] Decreased respiratory problems
 - [C] Loss of medicinal plants
 - [D] Increased soil erosion
23. Which of the following is an example of confounding in an epidemiological study?
- [A] A study on smoking and lung cancer that does not account for age
 - [B] Misclassification of exposure status in a case-control study
 - [C] Loss to follow-up in a cohort study
 - [D] Non-random allocation of participants in a clinical trial
24. Which type of epidemiological study is best for establishing causality?
- [A] Case-control study
 - [B] Cohort study
 - [C] Cross-sectional study
 - [D] Ecological study



25. An ecological study finds a high correlation between national rates of alcohol consumption and liver cirrhosis mortality. Assuming this relationship applies to all individuals within the population would be an example of

- [A] confounding bias
- [B] selection bias
- [C] ecological fallacy
- [D] information bias

26. Which of the following is an example of a cross-sectional study?

- [A] A study that follows a group of smokers and non-smokers over time to compare lung cancer incidence
- [B] A survey conducted in a population to assess the prevalence of hypertension at a single point in time
- [C] A case-control study investigating the risk factors for breast cancer
- [D] A clinical trial testing a new vaccine

27. As per the National Health Mission (NHM) norms in tribal, hilly and difficult to access areas, one Community Health Center (CHC) should be available for every

- [A] 50000 population
- [B] 80000 population
- [C] 100000 population
- [D] 120000 population



28. In India, Ministry concerned with Integrated Child Development Scheme is

- [A] Ministry of Labour and Human Rights
- [B] Ministry of Health and Family Welfare
- [C] Ministry of Women and Child Development
- [D] Drug Controller General of India

29. International Red Cross is based in

- [A] Geneva
- [B] New York
- [C] New Delhi
- [D] Rome

30. Odds Ratio is calculated in

- [A] case-control study
- [B] cohort study
- [C] ecological study
- [D] randomized controlled trial

31. The spread of infectious disease can be explained by

- [A] Epidemiological Wheel Theory
- [B] Epidemiological Triad
- [C] Web of Causation
- [D] Maslow's Hierarchy of Needs

32. Which of the following human diseases is caused by bacteria?

- [A] Hepatitis A
- [B] Typhoid
- [C] Measles
- [D] Polio

33. Medical Council of India has been renamed as

- [A] National Medical Commission
- [B] National Medical Council
- [C] Ayushman Bharat
- [D] Bharat Nirrog Samiti

34. National Vector Borne Disease Control Programme includes all, **except**

- [A] Malaria
- [B] Filaria
- [C] Japanese Encephalitis
- [D] Yellow Fever



35. Population of geriatric (>60 years) in the State of Arunachal Pradesh is

- [A] 5.2%
- [B] 10.3%
- [C] 15%
- [D] 8.8%

36. The full form of AEFI is

- [A] Adverse Effects Following Immunization
- [B] Adverse Events Following Immunization
- [C] Alternate Effects of Immunization
- [D] Advisory After Effective Immunization

37. As per World Health Organization, assistive products include

- [A] medication reminder
- [B] spectacles
- [C] crutch
- [D] All of the above

38. Birth Control Failure rate is measured as

- [A] Chandler Index
- [B] Pearl Index
- [C] Human Development Index
- [D] DMF Index

39. The infectious disease spreading rapidly with low fatality will have

- [A] high incidence and high prevalence
- [B] high incidence and low prevalence
- [C] low incidence and low prevalence
- [D] low incidence and high prevalence

40. The Pradhan Mantri Jan Arogya Yojana offers _____ to the people below poverty line.

- [A] health cover of ₹ 5 lakhs/family per year
- [B] disability cover of ₹ 5 lakhs/family per year
- [C] assured employment to youth less than 35 years of age
- [D] accidental cover to population

41. Which of the following is **correct** about forceps delivery?

- [A] Outlet forceps is applied when head is at or above 0 station
- [B] Can be applied with cervix not fully dilated
- [C] Risk of cephalohematoma is more than vacuum
- [D] OASI (Obstetric Anal Sphincter Injury) is more common than vacuum

42. HELLP syndrome is a complication of

- [A] transverse lie
- [B] pre-term labour
- [C] premature rupture of membranes
- [D] pre-eclampsia

43. Which of the following is **not true** about eclampsia?

- [A] 20–38% cases do not demonstrate hypertension or proteinuria before the seizure episode
- [B] Injection Levetiracetam is the drug of choice
- [C] Zuspan and Pritchard regimes are both effective in management
- [D] Fetal bradycardia following seizures usually recovers within few minutes

44. Magnesium sulfate is **not** indicated in which of the following obstetric situations?

- [A] Severe pre-eclampsia
- [B] Eclampsia
- [C] Fetal neuroprotection
- [D] GDM

45. Which of the following is **not true** about complete mole?

- [A] Histologically, it shows florid cistern formation, trophoblastic proliferation and absence of fetal parts
- [B] It has 46, XX chromosomes with both X chromosomes from paternal origin
- [C] There is no risk of progression to gestational trophoblastic neoplasia
- [D] Follow up with β HCG is required after suction and evacuation

46. Use of which of the following drugs is contraindicated in pregnancy?

- [A] Digoxin
- [B] Nifedipine
- [C] Amoxicillin
- [D] Enalapril



47. Methylergometrine is contraindicated for which of the following conditions?

- [A] Severe atonic uterine bleeding following delivery
- [B] Induction of labour
- [C] Uterine bleeding following abortion
- [D] Following expulsion of hydatidiform mole

48. Positive Fern test in cervical mucous smear is due to

- [A] the presence of NaCl under progesterone effect
- [B] the presence of NaCl under estrogen effect
- [C] LH/FSH
- [D] the mucus secretion by glands

49. Sonosalpingography is done
- [A] for measuring basal body temperature
 - [B] to detect pregnancy
 - [C] for testing tubal patency
 - [D] for determining an ovulatory cycle
50. Bacterial vaginosis is characterized by
- [A] fungal infection
 - [B] fishy vaginal odor
 - [C] vaginal pH < 4.5
 - [D] whiff test negative
51. Strawberry cervix is seen in
- [A] HPV infection
 - [B] trichomoniasis
 - [C] candidiasis
 - [D] cervical cancer
52. All of the following are **true** about uterine fibroid, **except**
- [A] estrogen and progesterone promote its development
 - [B] symptoms can include heavy menstrual bleeding, pain and heaviness
 - [C] red degeneration is seen during pregnancy
 - [D] sarcomatous change cannot occur
53. All of the following agents are used for emergency contraception, **except**
- [A] Cu-T
 - [B] Levonorgestrel
 - [C] high dose combined with oral contraceptives
 - [D] Misoprostol



54. Tubal sterilization is commonly performed at which site of fallopian tube?
- [A] Ampulla
 - [B] Infundibulum
 - [C] Isthmus
 - [D] Cornua
55. Postpartum IUCD can be inserted up to
- [A] 12 hours
 - [B] 24 hours
 - [C] 48 hours
 - [D] 72 hours
56. Which of the following is **not correct** about Cervical Intraepithelial Neoplasia (CIN)?
- [A] Mitotic activity, cellular immaturity, nuclear atypia in lower third of epithelium is CIN 1
 - [B] It typically originates inside the transformation zone
 - [C] Persistent high risk HPV infection is the main risk factor
 - [D] Coinfection with HIV does not impact its progression
57. All of the following are **correct** regarding tumour markers, **except**
- [A] CA-125-epithelial ovarian
 - [B] Inhibin B-trophoblastic tumour
 - [C] Alpha fetoprotein-yolk sac tumour
 - [D] LDH-dysgerminoma

58. Which of the following is **false** about Krukenberg tumour ovary?

- [A] Characterized by mucin-filled signet ring cells
- [B] Constitutes 30% to 40% of metastatic cancers to the ovaries
- [C] Primary tumour is frequently located in the stomach
- [D] They are usually unilateral solid masses

59. Regarding polycystic ovary syndrome in adolescents, which one is **true**?

- [A] Anti-Mullerian Hormone (AMH) levels assist in diagnosis
- [B] Glycemic status should be assessed like in adults
- [C] LH/FSH ratio is essential to make the diagnosis
- [D] Ultrasound should not be used to make the diagnosis

60. All of the following are **true** about cervical cancer, **except**

- [A] bimodal distribution of cases
- [B] it is mainly caused by HSV infection
- [C] vaccine decreases the incidence
- [D] it is an AIDS defining illness in HIV patients

61. Regarding HPV vaccine in India, which of the following is **false**?

- [A] Cervavac is indigenous vaccine of India
- [B] It is a bivalent vaccine against HPV 16 and 18
- [C] Approved between 9 and 26 years of age
- [D] It prevents anal cancer, genital warts along with cervical cancer

62. Presence of decubitus ulcer in prolapse indicates

- [A] infection
- [B] malignancy
- [C] circulatory disturbance
- [D] mechanical trauma



63. All of the following are signs of shoulder dystocia, **except**

- [A] turtle-neck sign
- [B] failure of restitution of fetal head
- [C] failure of shoulder to descend
- [D] difficulty with delivery of trunk

64. According to Anemia Mukh Bharat Program, 2018, strategies to decrease anemia in pregnancy includes all, **except**

- [A] daily, 1 iron and folic acid tablet starting from the fourth month of pregnancy
- [B] minimum 180 days during pregnancy and minimum 180 days postpartum
- [C] each tablet containing 100 mg elemental iron + 500 mcg folic acid
- [D] periodic deworming

65. Complications of Cesarean section includes all, **except**

- [A] puerperal infection
- [B] wound hematoma
- [C] venous thromboembolism
- [D] arterial thrombosis

66. Regarding screening for gestational diabetes mellitus, which of the following is **correct**?

- [A] Screening is done at 24 weeks only
- [B] OGTT according to DIPSI is done in fasting state
- [C] OGTT is done with 100 gm glucose
- [D] According to DIPSI criteria, 2 hours blood glucose >140 mg/dl is abnormal

67. All are **correct** regarding antenatal corticosteroid, **except**

- [A] it should be given between 24 to 34 + 6 weeks
- [B] it should be given when delivery is anticipated within 7 days
- [C] double the dose of steroids is required in twin pregnancy
- [D] diabetic women require additional dose of insulin while receiving steroids

68. The characteristics of caput succedaneum include all, **except**

- [A] crosses midline
- [B] results in neonatal jaundice
- [C] it disappear spontaneously within 4-6 days
- [D] it's a diffuse edematous swelling of the soft tissues of the scalp

69. Which is the best modality for determining gestational age when LMP is **not** reliable?

- [A] First trimester ultrasound
- [B] 26-28 weeks ultrasound
- [C] Date of Urine Pregnancy Test (UPT)
- [D] Uterine size by first trimester P/V exam

70. All of the following are **correct** about postmenopausal bleeding, **except**

- [A] endometrial atrophy accounts for 60% to 80% of cause
- [B] endometrial polyps account for 2% to 10% of cause
- [C] endometrial thickness >8 mm needs evaluation
- [D] hysteroscopy and endometrial biopsy are recommended

71. Which of the following is **not** a physiological change in pregnancy?

- [A] Increase in intravascular volume
- [B] Increase in cardiac output
- [C] Increase in stroke volume
- [D] Increase in peripheral vascular resistance

72. What is additional calories intake in pregnancy?

- [A] Less than 200 kcal
- [B] 200 kcal
- [C] 300 kcal
- [D] No change

73. All of the following increase in pregnancy, **except**

- [A] vital capacity
- [B] functional residual capacity
- [C] residual volume
- [D] tidal volume



74. Components of antenatal care is all, **except**

- [A] advise regarding antenatal visits
- [B] to make a plan for mode and place of delivery
- [C] family planning advise
- [D] screen for rare maternal diseases

75. Presence of 'show' and formation of 'bag of waters' are characteristics of

- [A] true labour
- [B] false labour
- [C] prelabour
- [D] PROM

76. Which is **not correct** about postpartum hemorrhage?

- [A] Blood loss \geq 1000 ml during Cesarean section
- [B] Blood loss \geq 500 ml during vaginal delivery
- [C] Blood loss accompanied by signs or symptoms of hypovolemia occurring within 24 hours after birth uterine atony accounts for 70% cases
- [D] Shock index cannot alert about severity of PPH

77. The opening of vagina, urethra and ducts of vestibular glands open into

- [A] labia majora
- [B] uterus
- [C] vestibule
- [D] None of the above

78. Which is associated with the least chance of ectopic pregnancy?

- [A] Hysteroscopic tubal occlusion
- [B] Copper IUCD
- [C] Combined OC pills
- [D] Tubal ligation

79. Medical management is likely to fail in ectopic pregnancy, if

- [A] the gestational sac is <3 cm
- [B] the duration of pregnancy is <5 weeks
- [C] the cardiac activity is present
- [D] the beta HCG <5000 mIU/ml

80. Which of the following is the most appropriate management in ruptured tubal pregnancy in hypovolemic shock?

- [A] Quick resuscitation followed by laparotomy and excision of the actively bleeding affected fallopian tube
- [B] Blood transfusion and resuscitation only after the clamps have been put to control the bleeding
- [C] Immediate autotransfusion of only fresh blood harvested from the peritoneal cavity
- [D] Excision of the affected tube and the ipsilateral ovary (salpingo-oophorectomy)



81. Fractional Inactivated Polio Vaccine is administered at all of the following ages in National Immunization Schedule, **except**

- [A] 6 weeks
- [B] 10 weeks
- [C] 14 weeks
- [D] 9 months

82. Which of the following activities is covered under the Rashtriya Bal Suraksha Karyakram?

- [A] Management of acute respiratory tract infection
- [B] Correction of dehydration in diarrhea
- [C] Screening for developmental delay
- [D] Catch-up immunization

83. Which of the following is a serious adverse event after immunization?

- [A] Local injection site pain
- [B] Permanent disability
- [C] Fever
- [D] Irritability



84. Deficiency of which of the following vitamins causes classical triad of diarrhea, dermatitis and dementia?

- [A] B12
- [B] Riboflavin
- [C] Niacin
- [D] Pyridoxine

85. Which of the following measures can be adopted for population screening of iodine deficiency?

- [A] Home salt iodine estimation
- [B] Soil iodine estimation
- [C] Urinary iodine excretion
- [D] Thyroid function test

86. Which of the following components accurately describes the composition of human breast milk?

- [A] Proteins 3.3 g/dL
- [B] Fats 4.2 g/dL
- [C] Carbohydrate 5 g/dL
- [D] Energy 100 kcal/dL

87. Which of the following conditions presents with metabolic alkalosis in children?

- [A] Renal tubular acidosis
- [B] Cerebral salt wasting
- [C] Bartter syndrome
- [D] Methanol poisoning

88. Which of the following drugs can be used to treat acute kidney injury and hyperkalemia?

- [A] Insulin and glucose
- [B] 3 percent saline
- [C] Chlorothiazide
- [D] Dopamine

89. Which of the following veins can be used for central venous access?

- [A] Internal jugular vein
- [B] Anterior cubital vein
- [C] Popliteal vein
- [D] Saphenous vein

90. The ratio of chest compressions to rescue breaths in resuscitation of children (>12 mo) is

- [A] 3 : 1
- [B] 10 : 2
- [C] 15 : 2
- [D] 30 : 2

91. Which of the following is a priority (not emergency) sign for triaging a sick child as per F-IMNCI?

- [A] Convulsion
- [B] Severe dehydration
- [C] Shock
- [D] Jaundice

92. Which of the following is a sign of severe dengue?

- [A] Clinical fluid accumulation
- [B] Serum aminotransferase (ALT) > 1000 U/L
- [C] Mucosal bleeds
- [D] Increase in hematocrit with decrease in platelets

93. A newborn who presents with cataract, sensorineural hearing loss and thrombocytopenia should be investigated for which congenital infection?

- [A] Syphilis
- [B] Cytomegalovirus
- [C] Mumps
- [D] Human Papillomavirus

94. Acyclovir therapy is not indicated in uncomplicated varicella infection in childhood as

- [A] it is costly
- [B] it does not prevent meningoencephalitis
- [C] the drug has marginal benefit
- [D] it is associated with side effects

95. Which of the following is an X-linked disease?

- [A] Friedreich's ataxia
- [B] Fabry's disease
- [C] Osteogenesis imperfecta
- [D] Cystic fibrosis

96. The appropriate approach to a neonate presenting with vaginal bleeding on day 4 of life is

- [A] administration of vitamin K
- [B] investigation for bleeding disorder
- [C] no specific therapy
- [D] administration of 10 ml/kg of fresh frozen plasma over 4 hours

97. The most common malignant neoplasm of infancy is

- [A] malignant teratoma
- [B] neuroblastoma
- [C] Wilms' tumour
- [D] hepatoblastoma

98. The important fatty acid present in breast milk which is important for growth is

- [A] docosahexaenoic acid
- [B] palmitic acid
- [C] linoleic acid
- [D] linolenic acid

99. Delayed Cord Clamping is recommended after

- [A] 30 seconds
- [B] 1 minute
- [C] 4 minutes
- [D] 5 minutes

100. Choose the **correct** response for growth interpretation of Indian children.

- [A] WHO growth charts are recommended for the use in children from 1-18 years of age
- [B] WHO growth charts are recommended for the use in children from 5-18 years of age
- [C] IAP growth charts are recommended for the use in children from 1-5 years of age
- [D] IAP growth charts are recommended for the use in children from 5-18 years of age

SPACE FOR ROUGH WORK



Faint, illegible text on the right side of the page, possibly bleed-through from the reverse side.

SPACE FOR ROUGH WORK



SEAL

SPACE FOR ROUGH WORK



SEAL



★★★



SEAL

GDMO-ALLOPATHY**PAPER-II****SET - A**

Q.No.	ANSWER KEY
1	D
2	A
3	A
4	A
5	B
6	C
7	A
8	A
9	B
10	B
11	C
12	D
13	D
14	B
15	D
16	C
17	B
18	C
19	A
20	B
21	A
22	B
23	A
24	B
25	C
26	B
27	B
28	C
29	A
30	A
31	B
32	B
33	A
34	D
35	D

36	B
37	D
38	B
39	A
40	A
41	D
42	D
43	B
44	D
45	C
46	D
47	B
48	B
49	C
50	B
51	B
52	D
53	D
54	C
55	C
56	D
57	B
58	D
59	D
60	B
61	B
62	C
63	D
64	C
65	D
66	D
67	B
68	B
69	A
70	C
71	D
72	C
73	B,C
74	D

75	A
76	D
77	C
78	C
79	C
80	A
81	B
82	D
83	B
84	D
85	C
86	B
87	C
88	A
89	A
90	C
91	D
92	B
93	B
94	C
95	B
96	C
97	B
98	A
99	B
100	D