

321005

COMBINED COMPETITIVE EXAMINATION (MAIN)

MATHEMATICS

Paper-I

Time : 3 Hours

Full Marks : 200

- Note : (1) The figures in the right-hand margin indicate full marks for the questions.  
(2) Attempt five questions in all.  
(3) Question No. 1 is compulsory.

1. Answer any ten questions from the following: 4×10=40

(a) Let  $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a linear transformation such that  $T(1, 0) = (1, -1)$  and  $T(0, 1) = (2, 3)$ . Find  $T(x, y)$  for any  $(x, y) \in \mathbb{R}^2$ . Also show that  $T$  is one-one and onto.

(b) A function  $f: \mathbb{R} \rightarrow \mathbb{R}$  is defined as  $f(x) = x^2 \sin \frac{1}{x}$  if  $x \neq 0$ , and  $f(0) = 0$ . Show that  $f'(x)$  exist for  $x \neq 0$  but is not continuous at  $x = 0$ .

(c) Show that  $\Gamma\left(\frac{1}{z}\right) = \sqrt{\pi}$ .

(d) Find  $a$  so that the points  $(a, 0, 3)$  and  $(0, -1, 0)$  are equidistant from the plane  $2x - 3y + z = 5$ .

(e) Find a solution  $\phi$  of the equation  $y'' - 2y' - 3y = 0$  if  $\phi(0) = 0$  and  $\phi'(0) = 1$ .

(f) If  $r(t) = t^2\mathbf{i} + t\mathbf{j} - t^3\mathbf{k}$  find  $\int_1^2 r \times \frac{d^2r}{dt^2} dt$

(g) As a result of leakage, an electrical capacitor discharges at a rate proportional to the charge. If the charge  $Q$  has the value  $Q_0$  at the time  $t = 0$ , find  $Q$  as a function of  $t$ .

- (h) Show that the matrix  $A = \begin{pmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{pmatrix}$  satisfies  $A^2 - 4A - 5I = 0$  where  $I$  is the identity matrix of order  $3 \times 3$ . Also find  $A^{-1}$ .
- (i) Show that the total energy is the sum of classical kinetic energy and rest mass energy.
- (j) Show that the central attraction is inversely proportional to  $r^2$  if the central orbit is  $\frac{l}{r} = 1 + e \cos \theta$  with pole as the centre.
- (k) Show that the least velocity with which a body can be projected to have a horizontal range  $R$  is  $\sqrt{gR}$  m/s and the greatest height attained is  $\frac{R}{4}$ .
- (l) A particle rests inside a hollow sphere of radius  $a$ . If the coefficient of friction is  $\frac{1}{\sqrt{3}}$ , find the height of the particle from the lowest point.

2. Answer any *eight* questions from the following : 5×8=40

- (a) Show that every square matrix can be expressed as a sum of a symmetric and a skew-symmetric matrix.
- (b) Show that  $\int_0^{\infty} e^{-ax} \frac{\sin x}{x} dx$ ,  $a \geq 0$  is convergent.
- (c) Solve the equation  $xyp^2 + (x^2 + y^2)p + xy = 0$ .
- (d) Show that  $a^x > x^a$  if  $x > a \geq e$ .
- (e) Find the matrix of the linear transformation  $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$  such that  $T(x, y, z) = (2x - y + 2z, y + z, 2x - 2y + z)$  with respect to the standard basis. Also, find its rank.
- (f) Show that the covariant derivative of the either of the fundamental tensors is zero.
- (g) A particle of mass  $m$  is acted upon by a force  $m\mu \left( x + \frac{a^4}{x^3} \right)$  towards the origin. If it starts from rest at a distance  $a$  show that it will arrive at the origin in time  $\frac{\pi}{4\sqrt{\mu}}$ .
- (h) A particle is projected on the inside of a smooth vertical circle of radius  $a$  from its lowest point with a velocity  $u$ . Show that the particle goes right round the circle if  $u^2 > 5ag$ .
- (i) Let  $f$  be a differentiable function for all values of  $x$ ,  $-\infty < x < \infty$ , such that  $f(-3) = -3$ ,  $f(3) = 3$  and  $|f'(x)| \leq 1$ . Show that  $f(0) = 0$ .

(j) Solve the equation  $y'' + 4y = \cos x$ .

3. Answer any **five** questions from the following :

8×5=40

(a) Let  $f: \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be a linear transformation such that  $f(\alpha_1) = 1, f(\alpha_2) = -1, f(\alpha_3) = 3$  where  $\alpha_1 = (1, 0, -1), \alpha_2 = (-1, 1, 1)$  and  $\alpha_3 = (1, 1, 0)$ . Find the value of  $f(a)$  for any element  $a = (a, b, c) \in \mathbb{R}^3$ .

(b) Find the equation of the plane through the point  $(2, 5, -8)$  and perpendicular to each of the planes  $2x - 3y + 4z + 1 = 0$  and  $4x + y - 2z + 6 = 0$ .

(c) Show that the line  $y = m(x + a) + \frac{a}{m}$  touches the parabola  $y^2 = 4a(x + a)$ .

(d) If  $f = (x + y + 1)\mathbf{i} + \mathbf{j} + (-x - y)\mathbf{k}$ , show that  $f \cdot \text{curl } f = 0$ .

(e) Find the stability of equilibrium of a vessel containing a liquid floating in a liquid.

(f) Show that for any value of  $x$ ,  $-\frac{1}{2} \leq \frac{x}{1+x^2} \leq \frac{1}{2}$ .

(g) Show that  $\frac{\partial g_{ij}}{\partial x^k} - \frac{\partial g_{jk}}{\partial x^i} = [jk, i] - [ij, k]$ .

4. Answer any **four** questions from the following :

10×4=40

(a) Find the eigenvalues and eigenvectors of the matrix  $\begin{bmatrix} 0 & -2 & -3 \\ -1 & 1 & -1 \\ 2 & 2 & 5 \end{bmatrix}$

(b) A heavy particle of weight  $W$ , attached to a fixed point by a light inextensible string, describes a circle in a vertical plane. The tension in the string has values  $mW$  and  $nW$  respectively when the particle is at the highest and the lowest point of its path. Show that  $n = m + 6$ .

(c) Reduce the equation  $3x^2 - 6xy - 5y^2 - 6x + 22 - 17 = 0$  to the standard form. Is the conic a central conic? If so, find the centre.

(d) Prove that the minimum value of  $\frac{(2x-1)(x-8)}{x^2-5x+4}$  is greater than its maximum value.

(e) If  $y = \tan^{-1} x$ , show that  $(1 + x^2) y_{n+1} + 2nxy_n + n(n-1) y_n = 0$ .

5. Answer any *two* questions from the following :

20×2=40

(a) (i) If  $I_n = \int_0^{\pi} (a \cos \theta + b \sin \theta)^2 d\theta$ , where  $n$  is a positive integer not less than 2, show that  $n I_n = ab(a^{n-2} + b^{n-2}) + (n-1)(a^2 + b^2) I_{n-2}$ .

(ii) Show that  $\text{curl} \left( \frac{a \times r}{r^2} \right) = -\frac{a}{r^3} + \frac{3r}{r^3} (a \cdot r)$

(b) (i) Prove that the lines in which the plane  $x + y + z = 0$  cuts the cone  $ayz + bzx + cxy = 0$  are at right angles if  $a + b + c = 0$ .

(ii) Find the directional derivative of  $\phi(x, y, z) = xy + yz + zx$  at the point  $(1, 2, 0)$  in the direction of  $i + 2j + 2k$ . Find in which direction is the directional derivative maximum? Find its value. Also find the unit normal and tangent plane to the surface  $xy + yz + zx = 2$  at the point  $(1, 2, 0)$ .

(c) If an area is bounded by two concentric semi-circles with their common bounding diameter in the free surface, prove that the depth of the centre of pressure is

$$\frac{3}{16} \pi \frac{(a+b)(a^2+b^2)}{a^2+b^2+ab} \text{ where } a \text{ and } b \text{ are the radii.}$$

6. Answer any *four* of the following :

10×4=40

(a) Show that the function  $\phi_1(x) = e^x$  is a solution of the differential equation

$$xy'' - (x+1)y' + y = 0.$$

Find a second independent solution of this differential equation.

(b) A rod of small section and of density  $\rho$ , has a small portion of metal of weight  $\frac{1}{n}$ th that of the rod attached to one extremity. Show that the rod will float at any inclination in a liquid of density  $\sigma$ , if  $(n+1)^2 \rho = n^2 \sigma$ .

(c) Let  $W_1, W_2, W_3$  be subspaces of a vector space. If  $W_2 \subseteq W_1$ , show that  $W_1 \cap W_2 + W_3 = W_2 + W_1 \cap W_3$ .

(d) Show that the dimension of the vector space of all real symmetric matrices of order  $n \times n$  is  $\frac{n(n+1)}{2}$ .

(e) Show that the equation of the straight line passing through the vector  $d$  and equally inclined to three mutually perpendicular vectors  $a, b, c$  is  $r = d + t \left( \frac{a}{|a|} + \frac{b}{|b|} + \frac{c}{|c|} \right)$  for  $t \in \mathbb{R}$ .

7. Answer any *two* questions from the following : 20×2=40

(a) Two particles  $A$  and  $B$  of mass  $m$  and one particle  $C$  of mass  $M$  are kept on the  $x$ -axis in the order  $A, B, C$ . Particle  $A$  is given a velocity  $vi$ . Consequently there are two collisions, both of which are completely inelastic. If the net energy loss because of these collisions is  $\frac{3}{4}$  of the initial energy, show that  $M = 2m$ .

(b) A particle of mass  $m$  is projected vertically under gravity, the resistance of the air being  $mk$  times the velocity. Show that the greatest height attained by the particle is  $\frac{V^2}{g} [\lambda - \log(1 + \lambda)]$ , where  $V$  is the terminal velocity of the particle and  $\lambda V$  is its initial velocity. Show that the corresponding time is  $\frac{V}{g} \log(1 + \lambda)$ .

(c) If the pressure of air varies as  $\left(1 + \frac{1}{m}\right)$ -th power of the density, show that, neglecting variation of temperature and gravity, the height of the atmosphere would be equal to  $(m+1)$  times the height of the homogeneous atmosphere.

8. Answer any *two* questions from the following : 20×2=40

(a) Six equal heavy uniform rods of weight  $w$  each are freely joined at their extremities. One rod is fixed in a horizontal position and the system lies in a vertical plane. The mid-points of the two upper non-horizontal rods are connected by a string. Show that the tension of the string is  $6w \cot \theta$  where  $\theta$  is the inclination of the non-horizontal rods to the horizontal.

(b) Two equal uniform rods are firmly joined at one end so that the angle between them is  $\alpha$  and they rest in a vertical plane on a smooth sphere of radius  $r$ . Show that they are in a stable equilibrium according as the length of the rod is  $>$  or  $< 4r \operatorname{cosec} \alpha$ .

9. Answer the following questions : 10+20+10=40

(a) Show that  $\lim_{n \rightarrow \infty} (\cos mx)^{\frac{n}{x^2}} = e^{\frac{1}{2}m^2n}$

(b) Let  $f(x, y) = x^2 \tan^{-1} \frac{y}{x} - y^2 \tan^{-1} \frac{x}{y}$  when  $x \neq 0$  and  $y \neq 0$  and  $f(x, 0) = f(0, y) = f(0, 0) = 0$ .

Show that  $f_{xy} = f_{yx}$  when  $x, y \neq 0$  but  $f_{xy}(0, 0) \neq f_{yx}(0, 0)$ .

(c) Find the center of gravity of the area enclosed by the curves  $y = mx$  and  $y^2 = 4ax$ .

10. Answer the following questions :

20×2=40

- (a) A particle of mass  $m$  is moving in +  $x$  direction with speed  $v$  and has momentum  $p$  and energy  $E$  in the frame  $S$ . If  $S'$  is moving at a speed  $v$  in the standard way and  $p'$  and  $E'$  are the momentum and energy respectively in  $S'$ , show that  $E'^2 - p'^2 c^2 = E^2 - p^2 c^2$ .
- (b) A spherical shell formed of two halves in contact along a vertical plane is filled with water. Show that the resultant pressure on either half of the shell is  $\frac{1}{4}\sqrt{13}$  of the total weight of the liquid.