

SEAL

DO NOT OPEN THE SEAL UNTIL INSTRUCTED TO DO SO

Series :

a

Question Booklet No. 320021

DT/08/24

PAPER—II

CHEMISTRY

Invigilator's Signature

Candidate's Signature

Time : 2 Hours

Maximum Marks : 100

ROLL NO.

--	--	--	--	--	--

INSTRUCTIONS FOR CANDIDATES

1. Immediately after the commencement of the Examination, candidate should check that this Booklet does **NOT** have any unprinted, torn or missing pages/SI. No. etc. If any defect is found, candidates should not write or mark anything on the OMR RESPONSE SHEET and immediately report it to the room Invigilator for replacement by a Complete Question Booklet.
2. Candidate should carefully read the instructions on the back of the OMR RESPONSE SHEET. They should **NOT** write Name, mark, make any stray marking or write anything irrelevant on either side of the OMR RESPONSE SHEET. Remarks other than the Answers and requisite details will be treated as revealing your identity and upon physical verification, if such remarks are found, the OMR RESPONSE SHEET will be invalidated and the candidature cancelled. No rough work should be done on the OMR RESPONSE SHEET. Rough work space provided in the Question Booklet can be used for the same.
3. Candidate should enter correct and complete digits of his/her Roll Number, Booklet Number and other details in the appropriate boxes and darken the corresponding bubbles in the OMR RESPONSE SHEET.
4. Candidate should **NOT** handle his/her OMR RESPONSE SHEET in such a manner as to mutilate, fold etc.
5. This Question Booklet contains **100** questions carrying 1 (one) mark each. Each question contains four responses. **Only one response/answer** for each question should be marked appropriately in the bubble on the OMR RESPONSE SHEET. If more than one response is marked, the answer will be considered wrong.
6. **Candidates are strictly prohibited to possess any book, notebook or loose paper, calculator, mobile phone, any electronic gadget, digital wristwatch etc., inside the Examination Hall, except his/her Unique ID, Admit Card and writing materials only.**
7. Immediately after the final bell, indicating the closure of the Examination, candidates should stop marking answers. Candidates should remain seated till the collection of OMR RESPONSE SHEET by the Invigilator. They will leave the Examination Hall after submission of OMR only after they are permitted by the Invigilator.
8. Violation of any of the above Rules will render the candidate liable to be expelled and disqualified from the Examination and according to the nature and gravity of his/her offence, he/she may be debarred from future Examinations and interviews to be conducted by the Commission and other such organizations.

NB : CANDIDATES ARE ALLOWED TO TAKE THIS QUESTION BOOKLET WITH THEM ONLY AFTER COMPLETION OF 2 (TWO) HOURS OF EXAMINATION TIME.

DO NOT OPEN THE SEAL UNTIL INSTRUCTED TO DO SO

/19-a

SEAL

SEAL

1. Which one of the following is a set of isoelectronic species?

[A] N_2, H_2S, CO

[B] N_2, CO_2, CN^-

[C] Ca, Mg, Cl

[D] N_2, CO, CN

2. ${}_{17}Cl^{35}$ and ${}_{17}Cl^{37}$ differ from each other in number of

[A] electrons

[B] protons

[C] nucleons

[D] mesons

3. The quantum number that is not obtained from the solution of Schrodinger's wave equation is

[A] principal quantum number

[B] azimuthal quantum number

[C] magnetic quantum number

[D] spin quantum number

4. An orbital with quantum numbers $n = 4,$

$l = 3, m = 0$ and $s = -\frac{1}{2}$ is called

[A] 4s orbital

[B] 4p orbital

[C] 4d orbital

[D] 4f orbital

5. The radius of hydrogen atom in the ground state is 0.53 \AA . The radius of Li^{2+} ion (atomic number = 3) in a similar state is

[A] 0.17 \AA

[B] 0.53 \AA

[C] 0.265 \AA

[D] 1.06 \AA



6. The outermost electronic configuration of the most electronegative element is

[A] ns^2np^3

[B] ns^2np^4

[C] ns^2np^5

[D] ns^2np^6

7. The first ionization potentials of Na, Mg, Al and Si are in the order

[A] $Na < Mg > Al < Si$

[B] $Na > Mg > Al > Si$

[C] $Na < Mg < Al > Si$

[D] $Na > Mg > Al < Si$

8. Which of the following has highest electron affinity?

[A] F

[B] Cl

[C] Br

[D] I

9. The general electronic configuration of transition elements is

- [A] $ns^2 (n-1) d^{10}$
- [B] $ns^{1-2} (n-1) d^{1-10}$
- [C] ns^1
- [D] $ns^2 np^5$

10. The shape and the type of hybridization about the central atom in $[I_3]^-$ are

- [A] trigonal planar, sp^2
- [B] pentagonal, sp^3d
- [C] linear, sp^3d
- [D] square pyramidal, d^2sp^3

11. The shape of sulphate ion is

- [A] tetrahedral
- [B] square planar
- [C] trigonal bipyramidal
- [D] hexagonal

12. Bond order is

- [A] directly related to bond length
- [B] inversely related to bond length
- [C] inversely related to bond strength
- [D] never fractional

13. The first important theory of coordination compounds was put forward by

- [A] Pauling
- [B] Slater
- [C] Werner
- [D] Lewis



14. The complex $CoCl_3 \cdot 5NH_3$ in aqueous solution ionizes to give a total number of chloride ions is equal to

- [A] 0
- [B] 1
- [C] 2
- [D] 3

15. The effective atomic number of iron in $[Fe(CN)_6]^{3-}$ is

- [A] 34
- [B] 35
- [C] 36
- [D] 37

16. In the spectrochemical series, the magnitude of the crystal field splitting is maximum for which ion?

- [A] Cl^-
- [B] F^-
- [C] NO_2^-
- [D] CN^-

17. The crystal field splitting energy for octahedral (Δ_o) and tetrahedral (Δ_t) complexes is related to

[A] $\Delta_t = \frac{4}{9}\Delta_o$

[B] $\Delta_t = \frac{1}{2}\Delta_o$

[C] $\Delta_o = 2\Delta_t$

[D] $\Delta_o = \frac{4}{9}\Delta_t$

18. The weight of iron which will be converted into its oxide (Fe_3O_4) by the action of 18 g of steam will be (atomic weight of Fe = 56 u)

[A] 21 g

[B] 42 g

[C] 84 g

[D] 168 g

19. In the reaction,



the equivalent weight of $\text{Na}_2\text{S}_2\text{O}_3$ (mol. wt. = M) is equal to

[A] M

[B] $M/2$

[C] $M/3$

[D] $M/4$

20. 5.6 liters of a gas at NTP is found to have a mass of 11 g. The molecular mass of the gas is

[A] 22

[B] 32

[C] 44

[D] 88

21. The free energy change for a reversible reaction at equilibrium is

[A] zero

[B] small positive

[C] small negative

[D] large positive

22. The solubility of CO_2 gas in water increases with

[A] increase in temperature

[B] reduction of gas pressure above water

[C] increase in volume

[D] increase of gas pressure above water

23. For a hypothetical reaction



occurring in a single step, the specific rate constants are 2.0×10^{-2} and 5.0×10^3 respectively for the forward and the backward reactions. The value of equilibrium constant is


[A] 4.0×10^{-4}

[B] 2.5×10^{-6}

[C] 2.5×10^5

[D] 4.0×10^{-6}

24. In the chemical reaction of gaseous reactants and product



increasing total pressure leads to

[A] increase in the amount of SO_3

[B] increase in the partial pressure of O_2


[C] increase in the partial pressure of SO_2

[D] change in the equilibrium constant



25. A solution of NH_4Cl is
- [A] acidic
 - [B] basic
 - [C] neutral
 - [D] amphoteric
26. Conjugate base of a strong acid is
- [A] a weak base
 - [B] a strong base
 - [C] a weak acid
 - [D] neutral
27. The pH value of 0.01 M HCl is equal to
- [A] 2
 - [B] 12
 - [C] 4
 - [D] 10
28. Which of the following is a weak electrolyte?
- [A] HCl
 - [B] NH_4Cl
 - [C] NaOH
 - [D] NH_4OH
29. Ostwald's dilution law is applicable to
- [A] strong electrolytes
 - [B] weak electrolytes
 - [C] non-electrolytes
 - [D] strong as well as weak electrolytes

30. One mole of N_2 and 3 moles of H_2 are mixed in 1.0 litre flask. If 50% N_2 is converted into ammonia by the reaction,



then the total number of moles of gas at equilibrium is

- [A] 1.5
 - [B] 3.0
 - [C] 4.5
 - [D] 6.0
31. Seaweeds are an important source of
- [A] chlorine
 - [B] bromine
 - [C] iodine
 - [D] zinc
32. In roasting, the ores are generally converted into
- [A] metals
 - [B] metal oxide
 - [C] hydrated metal oxides
 - [D] None of the above

33. The reducing agent used in thermite process is
- [A] magnesium
 - [B] chromium
 - [C] aluminium
 - [D] iron
34. Froth floatation process may be used to increase the concentration of the mineral in
- [A] bauxite
 - [B] chalcopyrites
 - [C] haematite
 - [D] calamine
35. White phosphorus contains
- [A] P₂ molecules
 - [B] P₄ molecules
 - [C] P₅ molecules
 - [D] P₆ molecules
36. Which of the following gases would have the highest RMS velocity at 25 °C?
- [A] Oxygen
 - [B] Carbon monoxide
 - [C] Carbon dioxide
 - [D] Sulphur dioxide
37. If the absolute temperature of a gas is doubled and the pressure is reduced to one-half, the volume of the gas will
- [A] be doubled
 - [B] remain unchanged
 - [C] increase four-fold
 - [D] be reduced to 1/4th
38. The unit of the van der Waals constant a is
- [A] atm L² mol⁻²
 - [B] atm L⁻² mol⁻²
 - [C] atm L mol⁻¹
 - [D] atm L⁻¹ mol
39. An ideal gas cannot be liquefied because
- [A] its critical temperature is always above 0 °C
 - [B] it solidifies before becoming a liquid
 - [C] forces operative between its molecules are negligible
 - [D] its molecules are relatively smaller in size



40. How many moles of He gas occupy 22.4 litres at 30 °C and one atmospheric pressure?

[A] 0.11

[B] 0.90

[C] 1.11

[D] 1.0

41. The rate of diffusion of oxygen as compared with ozone will be

[A] 0.66 times

[B] 0.82 times

[C] 1.22 times

[D] 1.5 times

42. At higher altitude the boiling point of water lowers because

[A] atmospheric pressure is low

[B] temperature is low

[C] atmospheric pressure is high

[D] None of the above

43. With rise in temperature, viscosity of a liquid

[A] increases

[B] decreases

[C] remains constant

[D] may increase or decrease

44. Among the following which crystal will be soft and have low melting point?

[A] Metallic

[B] Covalent

[C] Ionic

[D] Molecular

45. During evaporation of a liquid

[A] the temperature of the liquid rises

[B] the temperature of the liquid falls

[C] the temperature of the liquid unaffected

[D] None of the above

46. The kinetic energy of one mole of any gas depends upon

[A] pressure of the gas

[B] nature of the gas

[C] absolute temperature of the gas

[D] volume of the gas



47. In a solid lattice the cation has left a lattice site and is located at an interstitial position, the lattice defect is

- [A] vacancy defect
- [B] interstitial defect
- [C] Frenkel defect
- [D] Schottky defect



48. The second order Bragg diffraction of X-rays with $\lambda = 1.0 \text{ \AA}$ from a set of parallel planes in a metal occurs at an angle of 60° . The distance between the scattering planes in the crystal is

- [A] 0.575 \AA
- [B] 1.00 \AA
- [C] 1.15 \AA
- [D] 2.00 \AA


49. In crystal structure of sodium chloride, the arrangement of Cl-ions is

- [A] f.c.c.
- [B] b.c.c.
- [C] both f.c.c. and b.c.c.
- [D] None of the above

50. The hybridization of carbon in the structure of benzene is

- [A] sp
- [B] sp^2
- [C] sp^3
- [D] dsp^2

51. IUPAC name for the following compound is



- [A] *cis*-2-chloro-3-iodo-2-pentene
- [B] *trans*-2-chloro-3-iodopent-2-ene
- [C] *trans*-3-iodo-4-chloro-3-pentene
- [D] *cis*-3-iodo-4-chloro-3-pentene

52. Structures, CH_3COCH_3 and $\text{CH}_2=\text{CH}(\text{OH})\text{CH}_3$ represent

- [A] functional isomerism
- [B] metamerism
- [C] keto-enol tautomerism
- [D] position isomerism

53. A mixture of equal parts of (+) and (-) enantiomers is called

- [A] racemic mixture
- [B] homogeneous mixture
- [C] equilibrium mixture
- [D] resonance hybrid

54. An important chemical method to resolve a racemic mixture makes use of the formation of

- [A] a meso compound
- [B] enantiomers
- [C] diastereomers
- [D] racemates

55. A functional isomer of 1-butyne is

- [A] 2-butyne
- [B] 1-butene
- [C] 2-butene
- [D] 1,3-butadiene

56. Which of the following will show geometrical isomerism?

- [A] Propene
- [B] 1-butene
- [C] Isobutylene
- [D] 1,2-dibromobutane

57. On heating sodium phenoxide with methyl iodide, we get

- [A] anisole
- [B] diethyl ether
- [C] diphenyl ether
- [D] phenol



58. Glycerol is a

- [A] primary alcohol
- [B] monohydric alcohol
- [C] secondary alcohol
- [D] trihydric alcohol

59. Which one of the following compounds will be most readily attacked by an electrophile?

- [A] Benzene
- [B] Chlorobenzene
- [C] Toluene
- [D] Phenol


60. Aromatic aldehydes, in the presence of cyanide ion as catalyst, are converted into acylloins. This reaction is called

- [A] Perkin reaction
- [B] Benzoin condensation
- [C] Cannizzaro reaction
- [D] Claisen condensation

61. Which of the following compounds gives a ketone with Grignard's reagent?

- [A] Ethyl alcohol
- [B] Formaldehyde
- [C] Ethanenitrile
- [D] Methyl iodide

62. The product **B** in the following sequence of reactions is




- [A] acetone
[B] acetic acid
[C] isopropyl alcohol
[D] ethanol
63. Which of the following is least acidic?
- [A] $\text{C}_2\text{H}_5\text{OH}$
[B] $\text{C}_6\text{H}_5\text{OH}$
[C] CH_3COOH
[D] ClCH_2COOH
64. In presence of acid, hydrolysis of methyl cyanide gives
- [A] methylamine
[B] acetic acid
[C] methyl alcohol
[D] formic acid
65. The reaction between CO_2 and a Grignard reagent will yield
- [A] an alkene
[B] an alcohol
[C] a carboxylic acid
[D] an alkylmagnesium halide

66. Which of the following cannot reduce Fehling's solution?

- [A] Glucose
[B] Acetic acid
[C] Formaldehyde
[D] Acetaldehyde



67. In the following reaction,



the product **A** is

- [A] phenyl cyanide
[B] phenyl isocyanide
[C] ethylene chloride
[D] chlorobenzene
68. Which one of the following is a typical example of zwitterion?
- [A] Aniline
[B] Acetamide
[C] Aminophenol
[D] Glycine
69. The nature of glycosidic bond present in the structure of starch is
- [A] β - (1 \rightarrow 4)-glycosidic bond
[B] α - (1 \rightarrow 4)-glycosidic bond
[C] α - (1 \rightarrow 4)-glycosidic bond and α - (1 \rightarrow 6)-glycosidic bond
[D] None of the above

70. Which one of the following is **not** present in DNA?

[A] Adenine

[B] Cytosine

[C] Uracil

[D] Guanine



71. Adenosine consists of

[A] adenine and α -D-ribose

[B] adenine and β -D-ribose

[C] adenine and 2-deoxy-ribose

[D] None of the above

72. Which of the following is **not** present in nucleoside?

[A] Nucleobase

[B] Pentose sugar

[C] Phosphate

[D] None of the above

73. Sugar molecule present in the structure of RNA is

[A] α -D-ribose

[B] β -D-ribose

[C] β -D-deoxyribose

[D] glucose

74. Among the following, which is **not** optically active amino acid?

[A] Glycine

[B] Lysine

[C] Alanine

[D] Serine

75. The change in specific optical rotation of D-glucose in aqueous solution with time to an equilibrium value is called

[A] optical rotation

[B] mutarotation

[C] anomers

[D] None of the above

76. Which of the following is **not** a monosaccharide?

[A] Ribose

[B] Galactose

[C] Glycogen

[D] Fructose

77. The plant hormone responsible for fruit ripening is

[A] auxin

[B] cytokinin

[C] ethylene

[D] traumatic

78. The hormone that regulates retention of water in the kidney is

- [A] oxytocin
- [B] thyroxin
- [C] vasopressin
- [D] prolactin



79. Which of the following vitamins' deficiency causes scurvy?

- [A] Vitamin A
- [B] Vitamin C
- [C] Vitamin D
- [D] Vitamin K

80. Which of the following is also known as vitamin A?

- [A] Riboflavin
- [B] Retinol
- [C] Thiamine
- [D] Pyridoxin

81. The carbohydrate, which cannot be digested by the human digestive system, is

- [A] starch
- [B] glucose
- [C] fructose
- [D] cellulose

82. For a reversible isothermal process in equilibrium, the entropy change is given by the expression

[A] $\Delta S = \frac{T}{q_{\text{rev}}}$

[B] $\Delta S = \frac{q_{\text{rev}}}{T}$

[C] $\Delta S = \frac{\Delta V}{T}$

[D] $\Delta S = \frac{\Delta E}{T}$

83. Entropy change of a system and its surroundings in equilibrium

- [A] increases
- [B] decreases
- [C] remains constant
- [D] either increases or decreases

84. $\Delta H_{\text{combustion}}$ of a compound is

- [A] positive
- [B] negative
- [C] zero
- [D] positive or negative

85. Which law of thermodynamics introduces the concept of entropy?

- [A] First law
- [B] Second law
- [C] Zeroth law
- [D] Third law

86. Which among the following is an extensive property of the system?

- [A] Temperature
- [B] Refractive index
- [C] Volume
- [D] Viscosity



87. Heat exchanged in a chemical reaction at constant temperature and pressure is called

- [A] enthalpy
- [B] entropy
- [C] internal energy
- [D] free energy

88. The enthalpies of formation of $C_2H_4(g)$, $CO_2(g)$ and $H_2O(l)$ at $25^\circ C$ and 1 atm. pressure are 52 kJ mol^{-1} , -394 kJ mol^{-1} and -286 kJ mol^{-1} respectively. The enthalpy of combustion of $C_2H_4(g)$ will be

- [A] $+1412 \text{ kJ mol}^{-1}$
- [B] $-1412 \text{ kJ mol}^{-1}$
- [C] $+141.2 \text{ kJ mol}^{-1}$
- [D] $-141.2 \text{ kJ mol}^{-1}$

89. Energy required to dissociate 4 g of gaseous hydrogen into free gaseous atoms is 208 kcal at $25^\circ C$. The bond energy of H—H bond will be

- [A] 10.4 kcal
- [B] 104 kcal
- [C] 1040 kcal
- [D] 1.04 kcal

90. Two moles of an ideal gas expand spontaneously into a vacuum. The work done is

- [A] 1 J
- [B] 2 J
- [C] 4 J
- [D] zero

91. According to third law of thermodynamics, the entropy at $0^\circ K$ is zero for

- [A] elements in their stable form
- [B] perfectly crystalline solids
- [C] substances at 1 atm and $25^\circ C$
- [D] gaseous substances only

92. In which one of the following compounds the oxidation number of oxygen is positive?

- [A] H_2O_2
- [B] OF_2
- [C] Na_2O_2
- [D] H_2O

93. The oxidation number and covalency of sulphur in sulphur molecule (S_8) are

- [A] 0 and 2
- [B] +6 and 8
- [C] 0 and 6
- [D] +6 and 2

94. In the reaction,



the element which loses as well as gains electron is

- [A] Na
- [B] O
- [C] Cl
- [D] None of the above

95. The violent reaction between sodium and water is an example of

- [A] reduction
- [B] oxidation
- [C] redox reaction
- [D] neutralization reaction

96. In galvanic cell

- [A] oxidation occurs at cathode
- [B] oxidation occurs at anode
- [C] no reaction occurs at cathode
- [D] reduction occurs at anode

97. During the electrolysis of fused NaCl, which reaction occurs at anode?

- [A] Chloride ions are oxidised
- [B] Chloride ions are reduced
- [C] Sodium ions are oxidised
- [D] Sodium ions are reduced

98. The molar conductivity of weak electrolyte at infinite dilution can be obtained from

- [A] Ostwald's law
- [B] Kirchhoff's law
- [C] Kohlrausch's law
- [D] Faraday's law

99. When the cell reaction attains a state of equilibrium, the EMF of the cell is

- [A] zero
- [B] positive
- [C] negative
- [D] not definite

100. Calculate the EMF of the cell at 25 °C



if, $E_{\text{Cr}^{+3}/\text{Cr}}^\circ = -0.74\text{V}$ $E_{\text{Fe}^{+2}/\text{Fe}}^\circ = -0.45\text{V}$

- [A] +0.3 V
- [B] -0.3 V
- [C] +0.26 V
- [D] -0.26 V



SPACE FOR ROUGH WORK



SEAL

SPACE FOR ROUGH WORK

SEAL

/19-a

16



DDD24/4(171)—

SEAL